

Spontaneous Tuberculosis in Fishes and in Other Cold-blooded Vertebrates with Special Reference to *Mycobacterium fortuitum* Cruz from Fish and Human Lesions

ROSS F. NIGRELLI* & HENRY VOGEL*

New York Aquarium

(Plates I-VI)

INTRODUCTION

INTEREST in tuberculosis in cold-blooded animals was stimulated by a report by Bataillon, Dubard & Terre in 1897 on the disease in carp in a pond contaminated with dejecta from tubercular persons. It was later recognized that the acid-fast bacillus causing tuberculosis in the carp was a distinct species for which the name *Mycobacterium piscium* was given by Bataillon, Moeller & Terre (1902). The fact that this disease is found in fish led to speculations along the following lines: (1) that fish may be carriers of human tuberculosis organisms, (2) that fish, amphibians and reptiles could be used for the transmutation of human tuberculosis bacillus by serial passage, and (3) that these, or the acid-fast bacilli from cold-blooded animals, could be used for the treatment and prevention of human tuberculosis. The controversies engendered by discussions on these topics were reviewed by Vogel (1956, 1958) and by Parisot (1958).

The relatively recent discoveries of atypical human pathogenic species, *Mycobacterium fortuitum* Cruz (see Gordon, 1957) and *Mycobacterium balnei* Linell & Norden (1954), have revived interest in this subject.** These atypical forms, which are also found in fish and in water, together with the acid-fast bacilli causing tuberculosis in the lower vertebrates, show certain basic similarities in morphological, cultural and biochemical characteristics. Of particular interest is the report by Ross & Brancato (1959) that

one of the strains of acid-fast bacilli isolated in our laboratory (Nigrelli, 1953) from the Neon Tetra (*Hyphessobrycon innesi*) is the same as *Mycobacterium fortuitum*, a pathogenic species first isolated from human and cattle lesions in South America.

This paper deals with further information on the Neon Tetra strains of mycobacteria, tuberculosis in other fishes in the New York Aquarium and with reports of the disease in fishes, amphibians and reptiles by other investigators.

TUBERCULOSIS IN FISHES IN THE NEW YORK AQUARIUM

Routine examinations of fish in the New York Aquarium and a search of the literature show that tuberculosis in poikilothermic animals is much more prevalent than is generally suspected. The host species are listed in Table IV. Table I lists those host species that were found infected in the New York Aquarium and from which the acid-fast bacilli were isolated and cultured.

PATHOLOGY OF TUBERCULOSIS IN FISHES

Tubercular lesions are found in gills, skin, muscle, heart, kidneys, spleen, liver, pancreas, mesenteries, gonads, eyes and brain. Lepromatous-like macular and necrotic skin and fin lesions are characteristic of tuberculosis in the Three-spot or Blue Gourami (Pl. V, fig. 9). The disease in the Neon Tetra is recognized externally by yellowish discoloration of the usually brilliant red markings on both sides of the hind part of the body (Pls. I & II, figs. 1, 2, 3). In most species there is no external evidence but the disease is recognized internally by the presence of extensive, yellowish adhesions or by numerous miliary-like tubercle bodies in various organs (Pls. II, III, V & VI, figs. 4, 5, 10, 11).

*Bureau of Laboratories, New York City Department of Health.

**See also Clark, H. Fred, & Charles C. Shepard, "Effect of Environmental Temperature on Infection with *Microbacterium marinum* (*balnei*) of Mice and a Number of Poikilothermic Species. *Jour. Bact.*, 86 (5): 1059-1069. Nov., 1963.

TABLE I. HOSTS FROM WHICH ACID-FAST BACILLI HAVE BEEN ISOLATED
(Cultures maintained at the New York Aquarium)

| Species | Common Name | Habitat |
|----------------------------------|----------------------------|------------------------|
| Tropical Freshwater spp. | | |
| <i>Hyphessobrycon innesi</i> | Neon Tetra | Peruvian Amazon |
| <i>Trichogaster trichopterus</i> | Three-spot or Blue Gourami | Tropical Far East |
| <i>Toxotes jaculator</i> | Archerfish | Philippines & Far East |
| Tropical Marine spp. | | |
| <i>Plectorhynchus</i> sp. | Sweet-lip | Pacific coral reefs |
| <i>Premnas aculeatus</i> | Spiny Clownfish | Pacific coral reefs |
| <i>Amphiprion percula</i> | Common Clownfish | Pacific coral reefs |
| <i>Amphiprion akallaopsis</i> | Skunk Clownfish | Pacific coral reefs |
| <i>Amphiprion xanthurus</i> | Chocolate Clownfish | Pacific coral reefs |
| <i>Amphiprion latilavivus</i> | White-saddle Clownfish | Pacific coral reefs |

In the Climbing Perch and in the Goldfish the lesions appear as numerous pearl-like bodies in the liver, kidneys and mesenteries (Pls. V & VI, figs. 10, 11). Emaciation, exophthalmia, lordosis and other body abnormalities may or may not be associated with the disease.

Histopathologically, tuberculosis in fishes resembles the tubercular picture in warm-blooded animals but with the following differences: milder or no inflammatory reactions, greater fibrous development, absence of typical giant cells and little or no caseation (Pls. II & IV, figs. 3, 4, 7, 8). The term "hard tubercle" (Pls. V & VI, figs. 10, 11) appears appropriate for the lesions in the Climbing Perch, Goldfish and several other species. These tubercles are formed by coalescence of several smaller units (Pl. VI, fig. 11). In some freshwater fishes some degree of caseation may occur but typically the tissue reaction consists of loosely organized masses of semi-necrotic cells with groups of acid-fast bacilli in the core of epitheloid-like elements. (Pls. II & IV, figs. 3, 7, 8).

The term "mycobacteriosis" was suggested by Parisot & Wood (1960) as being more appropriate for tuberculosis in fish. The suggestion was based on the absence of typical inflammatory responses to the infection in salmonoid fishes. However, since this is not true for fishes generally and since typical tubercles and other classical tissues reactions are present, tuberculosis is a valid term for the disease in fish as well as in other cold-blooded animals.

TRANSMISSION

Tuberculosis in cold-blooded vertebrates can be induced by parenteral injections of mycobacterial suspensions. However, it is generally conceded that the natural mode of infection is by ingestion of the organisms directly from the water, by eating infected tissues or contaminated

feed. Such origins of the infective organisms have been experimentally demonstrated for tuberculosis in tadpoles (Nonidez & Kahn, 1934, 1937)¹, in the Mexican platyfish (Baker & Hagen, 1942) and in snails (Michelson, 1961)². The increase in incidence of tuberculosis in hatchery-maintained salmon and trout is directly related to the increased usage of infected salmon carcass as feed for young salmon, a detrimental hatchery practice of which fishery biologists were unaware until recently (Wood & Ordal, 1958). The possibility of transovarian transmission or of the spread of the infection by contamination of sperms and eggs during stripping of mature stock fish or the spawning run fish was also considered. Although the results on the salmon studies were inconclusive (Ross & Johnson, 1962), our observations of tuberculosis in embryonic platyfish and guppies certainly suggest transovarian transmission as a definite possibility. In addition, the entry of infective organisms through lesions of the skin and gills caused by parasites or by mechanical injury should also be considered.

There is no information on the mechanism of spread of tuberculosis in fish. If the portal of entry is through lesions in the skin, the route must be through the lymphatic system or through the blood stream. Since the preponderance of evidence indicates that the infection is brought about by ingestion, the spread must then take place through the gastro-intestinal tract. Just how this is accomplished has not been determined, especially since we have not observed tubercular lesions in this organ (Pl. II, fig. 4).

¹Jose F. Nonidez & Morton C. Kahn. Tuberculosis induced in the tadpole by feeding. Proc. Soc. Exp. Biol. & Med., 31: 783. Experimental tuberculosis infection in the tadpole and the mechanism of spread. Amer. Rev. Tuberc., 36: 191.

²Edward H. Michelson. An acid-fast pathogen of fresh water snails. Amer. J. Trop. Med. & Hyg., 60: 423.

ISOLATION AND CULTURAL PROCEDURES

Isolation of the acid-fast bacilli from pathological materials in fishes is relatively simple. In the Neon Tetra, for example, the disease is readily recognized by the yellowish discoloration of the red markings. The fish is anaesthetized with urethane (methyl carbamate) or with MS-222 (tricaine methanesulphonate, Sandoz), and the lateral body wall in front of the vent is cauterized with a red-hot scalpel; the flesh adhering to the blade results in a raised flap, exposing the body cavity. A sterile bacteriological needle is plunged into the lesions, preferably into those in the kidneys, and the material is then transferred to slants of media ordinarily used to culture mycobacteria. In our laboratory, slants of Dorset's or Petroff's egg agar is used successfully. It is our experience that with this simple technique pure cultures of acid-fast bacilli are usually obtained. If difficulty is encountered, however, a portion of the disease tissue is digested with 4% sodium hydroxide for 15-30 minutes to destroy the contaminating organisms. The digest is neutralized with HCl and centrifuged. The sediment is then inoculated on one of the media suggested above. The digestion time with NaOH can be varied, depending on the sensitivity of the bacilli and the quantity of tissue to be digested. Diseased tissue can also be treated with trisodium phosphate, oxalic acid or with 6% sulfuric acid. Other techniques and media for handling tubercular fish tissue are suggested by Westgard (1959), and those found in any standard reference on methods in pathology and bacteriology may also be used. Since contamination of fish material may reach a high level in a short time, it is suggested that moribund fish be sacrificed and used for the isolation procedures.

SOME CHARACTERISTICS OF MYCOBACTERIA ISOLATED FROM FISHES IN THE NEW YORK AQUARIUM

The mycobacteria are easily demonstrated in smears of the skin and organ lesions by the Ziehl-Neelsen staining method (Pls. I & III, figs. 2, 5). The bacilli, which may or may not be seen within macrophages, are pleomorphic slender rods varying in length from 3 to 7 microns and show the bead-like constituents in the cell when stained, or when seen in electron microscope preparations (Pl. III, fig. 6). The organisms grow slowly as raised colonies on glycerol and egg agar slants when kept at room temperature. However, once growth is established, usually within one to three weeks, subcultures will grow more rapidly. Mycobacteria from tropical fishes

grow well at 28°C. and subcultures become luxuriant even at 37°C. (Pl. VI, fig. 12) The degree of pigment production, which may develop either in light or in the dark, varies with the strain and age of the culture. The mycobacteria isolated from freshwater tropical fishes grown on Dorset's or Petroff's media vary in color from cream to yellowish-green; bacilli from stenohaline tropical fishes vary from light to bright yellow color.

CLASSIFICATION OF MYCOBACTERIA OF COLD-BLOODED VERTEBRATES

Table II list the species of mycobacteria from fishes, amphibians and reptiles that have been studied in detail; some of these are recognized as valid species and are included in Bergey's Manual of Determinative Bacteriology.

The growth and nutritional requirements, antigenic structure, pathogenicity, source and habitat of *Mycobacterium piscium*, *Myco. marinum*, *Myco. ranae*, *Myco. thamnopheos* and *Myco. friedmanni* are summarized by Reed (1948). Gordon (1957) recognizes and characterizes only *Myco. marinum*. *Myco. platypocilus*, *Myco. thamnopheos* and *Myco. fortuitum*. The type of culture of *Mycobacterium piscium* is apparently lost while *Myco. ranae* and *Myco. friedmanni*, together with certain fish strains (e.g., from Halibut and Halibut roe), are considered to be identical with *Myco. smegmatis* and/or with *Myco. fortuitum* (Gordon & Smith, 1955). The information on *Mycobacterium anabanti* Besse (1949a) and *Myco. salmoniphilum* Ross (1960) was not available at the time to Gordon and her co-workers for evaluation. However, Gordon & Mihm (1959) reported that certain of the strains from trout and salmon are identical with *Myco. fortuitum*.

Except for the Neon Tetra strains, the mycobacteria of the fish listed in Table I have not been further characterized. Ross & Brancato (1959) considered the Neon Tetra strain 9-21H, which is a subculture of our H-strain, to be identical with *Mycobacterium fortuitum*. Some differences in the utilization of several substances as carbon source are noted between strain 9-21H and strains H and N as analyzed by Vogel (1959). These are shown in Table III and compared with *Mycobacterium marinum*, *Myco. fortuitum* from mammals and with the several strains of *Myco. salmoniphilum* which were recognized as *Myco. fortuitum* by Gordon and Mihm (1959).

DISCUSSION

As pointed out by Fregnan, Smith & Randall (1961), a great deal of attention has been

TABLE II. SPECIES OF MYCOBACTERIA OF POIKILOOTHERMS

| Mycobacteria | Host |
|--|---|
| | I. Fishes |
| <i>Mycobacterium piscium</i> (see Reed, 1948) | <i>Cyprinus carpio</i> , European Carp |
| <i>Mycobacterium marinum</i> Aronson, 1926 | <i>Abudefduf mauritii</i> , Sergeant Major <i>Micropogon undulatus</i> , Croaker <i>Centropristis striatus</i> , Sea Bass (Philadelphia Aquarium) |
| <i>Mycobacterium platypoecilus</i> Baker & Hagen, 1942 | <i>Platyzoecilus maculatus</i> , Mexican Platyfish (Cornell Univ.) |
| <i>Mycobacterium anabanti</i> Besse, 1949a | <i>Macropodus opercularis</i> , Paradisefish (France) |
| <i>Mycobacterium fortuitum</i> Cruz (see Gordon, 1957) Neon Tetra Strain 9-21 H (Ross & Brancato, 1959) | <i>Hyphessobrycon innesi</i> , Neon Tetra (New York Aquarium) |
| <i>Mycobacterium salmoniphilum</i> Ross, 1960 | <i>Oncorhynchus tshawytscha</i> , Chinook Salmon <i>Salmon gairdneri</i> , Steelhead Trout (Hatcheries, Oregon & Washington) |
| | II. Amphibians |
| <i>Mycobacterium ranae</i> (Küster, 1905) (see Reed, 1948) | European Frogs |
| | III. Reptiles |
| <i>Mycobacterium friedmanni</i> Holland, 1920 (see Reed, 1948) | <i>Chelone corticata</i> , Loggerhead Turtle (European Zoo) |
| <i>Mycobacterium thamnopheos</i> Aronson, 1929 | <i>Thamnophis sirtalis</i> , Garter Snake (U.S.A.) |

directed towards the difficult problem of differentiation and classification of saprophytic and pathogenic mycobacteria ever since the discovery of the tubercle bacillus by Robert Koch in 1882. No entirely satisfactory method has yet been developed and all attempts so far have led to considerable confusion. This is especially true for the attempts to classify the mycobacteria causing tuberculosis in fishes, amphibians and reptiles. For example, Gordon & Smith (1955) and Gordon & Mihm (1959) reported that several strains of mycobacteria from cold-blooded vertebrates were identical either with *Mycobacterium smegmatis* or *Myco. fortuitum*. Gordon (1957) considers *Myco. marinum*, *Myco. platypoecilus* and *Myco. thamnopheos* to be valid species. However, the discovery of *Mycobacterium balnei* in swimming pools and in human lesions (Linell & Norden, 1954; Swift & Cohen, 1962) has added to the confusion. Bojalil (1959) considers this species to be the same as *Myco. marinum* while McMillen & Kushner (1959) report that *Myco. marinum*, *Myco. platypoecilus* and *Myco. balnei* represent a single species which, by priority, should be *Myco. marinum* Aronson. In addition, McMillen (1960) indicates that *Myco. fortuitum* is a mutation or adaptation of *Myco. marinum*.

If the above reports are valid, then all strains of mycobacteria from marine and freshwater fishes, regardless of ecological and other biological factors, belong to either *Mycobacterium*

marinum originally described from Atlantic Coast fishes or to *Mycobacterium fortuitum* first reported from cattle and human abscesses. It is difficult to believe that either species is a ubiquitous pathogen of fish with no host specificity.

In any event, it is quite apparent that tuberculosis in cold-blooded animals, and especially in fishes, is much more widespread than is generally suspected. Although tuberculosis has been found mainly in fishes kept in aquaria, hatcheries and in fish holding ponds, cases of tuberculosis in feral populations have been reported (Nigrelli, 1953). Of particular interest is the epizootics in salmonoids in the Pacific Northwest, both in hatchery-reared fishes and in migrating populations (Wood & Ordal, 1958; Wood, 1959; Ross, 1959; Parisot & Wood, 1960). Tuberculosis in these species (Table IV), the presence of which was first reported by Earp, Ellis & Ordal (1955), is therefore of great economic importance. The disease is also prevalent in a large variety of tropical fresh water fishes. The members of the families Characidae and Cyprinidae, which are highly valued by fish hobbyists, are especially susceptible and commercially available to the microbiologist interested in this problem.

SUMMARY

A survey of fishes in the New York Aquarium and a search of the literature show that tuberculosis in these animals and in other poikilotherms is more prevalent than is generally suspected.

TABLE III. CARBON SOURCES UTILIZED BY MYCOBACTERIA FROM FISH

| Carbon Source | Neon Tetra Strains ¹ | | | | <i>Myc. fortuitum</i> ² | | Myc. marinum ³ VII | <i>Myc. salmoniphilum</i> ⁴ | | | | |
|---------------|---------------------------------|----------|-----|--------------|------------------------------------|----------|----------------------------------|--|----|---|----|---|
| | I | II | III | IV | V | VI | | VIII | IX | X | XI | |
| Glucose | - | - | x | x | x | - | - | - | - | - | - | - |
| Mannose | x | x | x | x | x | x | - | x | x | x | x | x |
| Trehalose | x | x | x | x | x | x | - | x | x | x | x | x |
| Fructose | x | x | x | - | x | x | x | x | 0 | 0 | 0 | 0 |
| Mannitol | 0 | x | 0 | (occasional) | 0 | 0 | - | x | 0 | 0 | 0 | 0 |
| Xylose | (slight) | 0 | 0 | (occasional) | 0 | 0 | - | 0 | 0 | 0 | 0 | 0 |
| Sorbitol | x | x | 0 | (rarely) | 0 | (slight) | 0 | 0 | 0 | 0 | 0 | 0 |
| Sucrose | x | 0 | 0 | - | 0 | 0 | - | 0 | 0 | 0 | 0 | 0 |
| Rhamnose | 0 | 0 | 0 | 0 | 0 | 0 | - | 0 | 0 | 0 | 0 | 0 |
| Dextrose | x | x | - | - | - | x | - | x | x | x | x | x |
| Dulcitol | 0 | 0 | 0 | 0 | 0 | 0 | - | 0 | 0 | 0 | 0 | 0 |
| Arabinose | 0 | (slight) | 0 | (occasional) | 0 | 0 | x | 0 | 0 | 0 | 0 | 0 |
| Lactose | 0 | (slight) | 0 | 0 | 0 | 0 | - | 0 | 0 | 0 | 0 | 0 |
| Galactose | - | - | 0 | (occasional) | 0 | - | 0 | 0 | 0 | 0 | 0 | 0 |
| Raffinose | (slight) | (slight) | 0 | (rarely) | 0 | 0 | - | 0 | 0 | 0 | 0 | 0 |
| Inositol | 0 | 0 | 0 | (occasional) | 0 | 0 | - | 0 | 0 | 0 | 0 | 0 |
| Maltose | (slight) | x | 0 | (rarely) | 0 | (slight) | - | 0 | 0 | 0 | 0 | 0 |
| Inulin | 0 | 0 | - | - | - | 0 | - | - | - | - | - | - |
| Erythritol | 0 | 0 | - | (occasional) | - | 0 | - | - | - | - | - | - |
| Amygdalin | (slight) | (slight) | - | - | - | (slight) | - | - | - | - | - | - |
| Salicin | x | 0 | - | - | - | 0 | - | - | - | - | - | - |
| Starch | - | - | - | (hydrolyzed) | - | - | - | - | - | - | - | - |

¹I, H-strain, II, N-strain, Vogel (1959); III, 9-21H strain, Ross & Brancato (1959).²IV, Gordon (1957); V, Ross & Brancato (1959).³VI, Vogel (1959); VII, Gordon (1957) (Gordon & Mihm (1959), 80 strains examined).⁴VIII, CC-strain from steelhead trout in Chamber Creek Hatchery; IX, OR-strain from steelhead trout in Oak Ridge Hatchery; X, CAR-strain from chinook salmon in Carson Hatchery; XI, SC-strain from Chinook salmon in Spring Creek Hatchery, Ross (1960) (see also Gordon & Mihm, 1959).

TABLE IV. FISHES, AMPHIBIANS AND REPTILES IN WHICH TUBERCULOSIS HAS BEEN REPORTED

| Species | Common Name | Reference |
|---|--------------------------------------|---|
| I. Teleostomi | | |
| 1. Salmonidae | | |
| (1) <i>Oncorhynchus gorbuscha</i> | Humpback Salmon | 47, 64 |
| (2) <i>Oncorhynchus keta</i> | Dog Salmon | 47 |
| (3) <i>Oncorhynchus kisutch</i> | Silver Salmon | 47, 64 |
| (4) <i>Oncorhynchus nerka</i> | Blueback Salmon | 47, 64 |
| (5) <i>Oncorhynchus tshawytscha</i> | Chinook or King Salmon | 15, 41, 47 |
| (6) <i>Salmo gairdneri</i> | Steelhead, Rainbow Trout | 47, 64 |
| 2. Osmeridae | | |
| (7) <i>Osmerus mordax</i> | American Smelt | 62 |
| 3. Umbridae | | |
| (8) <i>Umbra pygmaea</i> | Mud-Minnow | 51 |
| 4. Characidae | | |
| (9) <i>Aphyochorax rubripinnis</i> (<i>Aphiocharax rubrepinis</i>) | Bloodfin | NYA, 11, 27 |
| (10) <i>Gymnocorymbus ternetzi</i> | Black Tetra | NYA, 11, 27, 32 |
| (11) <i>Hemigrammus rhodostomus</i> | Rummy-Nosed Characin | NYA, 32 |
| (12) <i>Hemigrammus unilineatus</i> | Feather Fin | 11 |
| (13) <i>Hemigrammus erythrozona</i> (<i>Hyphessobrycon gracilis</i>) | Glowlight Tetra | 11 |
| (14) <i>Hemigrammus ocellifer</i> | Head-and-Tail Light | 2 |
| (15) <i>Hyphessobrycon bifasciatus</i> | Yellow Tetra | 27 |
| (16) <i>Hyphessobrycon flammeus</i> | Flame Characin | NYA, 11 |
| (17) <i>Hyphessobrycon cardinalis</i> | Cardinal Tetra | NYA, 11 |
| (18) <i>Hyphessobrycon innesi</i> | Neon Tetra | NYA, 11, 38, 39, (also Conroy 1963)* |
| (19) <i>Hyphessobrycon pulcher</i> | | 32 |
| (20) <i>Hyphessobrycon rosaeus</i> (= <i>ornatus</i>) | Rosy Barb | 32 |
| (21) <i>Hyphessobrycon callistus</i> (= <i>serpae</i>) | Serpa Tetra | NYA, 2, 27, 32 |
| (22) <i>Moenkhausia pittieri</i> | Pittier's Moenkhausia | NYA, 32 |
| (23) <i>Pyrhulina rachoviana</i> | Rachow's Pyrrhulian | 27 |
| (24) <i>Pristella riddlei</i> | Riddle's Pristella | 2 |
| 5. Cyprinidae | | |
| (25) <i>Notemigonus crysoleucas</i> | Dace or Roach | 62 |
| (26) <i>Barbus fluviatilis</i> | European Barb | 51 |
| (27) <i>Brachydanio albolineatus</i> (<i>Danio albolineatus</i>) | Pearl Danio | NYA, 11, 27, 38 |
| (28) <i>Brachydanio analipunctatus</i> | Tail-Spot Danio | NYA, 38 |
| (29) <i>Brachydanio nigrofasciatus</i> | Spotted Danio | NYA |
| (30) <i>Brachydanio rerio</i> (<i>Danio rerio</i>) | Zebra Danio | NYA, 11, 32, 38 |
| (31) <i>Danio malabaricus</i> | Giant Danio | NYA, 11, 27, 32 |
| (32) <i>Carassius auratus</i> (<i>C. oratus</i>) | Goldfish | 11, 27, 32 |
| (33) <i>Carassius carassius</i> | Crusian Carp | 32 |
| (34) <i>Cyprinus carpio</i> <i>Cyprinus carpio</i> | Common Carp Hi-Goi or Golden Carp | 7, 8, 32 23 |
| (35) <i>Puntius conchoniis</i> (<i>Barbus conchoniis</i>) | Rosy Barb | NYA, 11 |
| (36) <i>Puntius lineatus</i> (<i>Barbus lineatus</i>) | Lined Barb | 32 |
| (37) <i>Puntius nigrofasciatus</i> (<i>Barbus nigrofasciatus</i>) | Black Ruby Barb | 32 |
| (38) <i>Puntius phutunio</i> (<i>Barbus phutunio</i>) | Dwarf or Pygmy Barb | 32 |
| (39) <i>Puntius semifasciolatus</i> | Chinese Barb | NYA |

*David W. Conroy, Univ. Nac. de Buenos Aires, Lab. de Bact., Inst. de Biol. Marino; Microbid. Españ., 16: 47-54.

TABLE IV. FISHES, AMPHIBIANS AND REPTILES IN WHICH TUBERCULOSIS HAS BEEN REPORTED (*Continued*)

| Species | Common Name | Reference |
|--|---------------------------|---|
| (40) <i>Puntius tetrazona</i> (<i>Barbus sumatranus</i>) | Sumatra Barb | NYA, 32 |
| (41) <i>Puntius partipentazona</i> | Five-Banded Barb | 32 |
| (42) <i>Puntius ticto</i> (<i>Barbus ticto</i>) | Ticto Barb | 32 |
| (43) <i>Rasbora einthoveni</i> | Brilliant Rasbora | 32 |
| (44) <i>Rasbora heteromorpha</i> | Red Rasbora | NYA, 11 |
| (45) <i>Rasbora lateristriata</i> | Striped Rasbora | NYA |
| (46) <i>Rasbora leptosoma</i> | Slender-Bodied Rasbora | 32 |
| (47) <i>Rasbora trilineata</i> | Scissors-Tail | NYA |
| (48) <i>Tanichthys albonubes</i> | White-Cloud Mountain Fish | NYA, 38 |
| (49) <i>Idus melanotus</i> | Golden Orf | 32 |
| (50) <i>Tinca tinca</i> | Tench | 10 |
| 6. Siluridae | | |
| (51) <i>Silurus glanis</i> | Wels or European Catfish | 23 |
| 7. Bagridae | | |
| (52) <i>Rhamdia sapo</i> | | 32 |
| 8. Clariidae | | |
| (53) <i>Clarias dumerili</i> | | 27, 32 |
| 9. Loricariidae | | |
| (54) <i>Plectostomus punctatus</i> | | 32 |
| 10. Gadidae | | |
| (55) <i>Gadus collarias</i> | Atlantic Codfish | 1, 28 |
| 11. Cyprinodontidae | | |
| (56) <i>Aphyosemion australe</i> | Lyretail | 11 |
| (57) <i>Oryzias latipes</i> (<i>Aplocheilus latipes</i>) | Medaka | 10, 11 |
| (58) <i>Aplocheilus panchax</i> (<i>Panchax panchax</i>) | Panchax | 10 |
| (59) <i>Rivulus cylindraceus</i> | Cuban Rivulus | 11 |
| (60) <i>Cynolebias wolterstorffi</i> | (So. American Annals) | Roy L. Walford Aug. 1963† (Personal Communication) |
| (61) <i>Cynolebias adeoffi</i> | | Roy L. Walford Aug. 1963† (Personal Communication) |
| (62) <i>Cynolebias elongatus</i> | | Roy L. Walford Aug. 1963† (Personal Communication) |
| 12. Poeciliidae | | |
| (63) <i>Lebistes reticulatus</i> | Guppy | NYA, 11 |
| (64) <i>Xiphophorus helleri</i> | Swordtail | NYA, 11, 27, 32 |
| (65) <i>Platypoecilus maculatus</i> | Platyfish | NYA, 6, 11, 38 |
| (65a) <i>Mollisia sphenops</i> | Mollie | 43 |
| 13. Belonidae | | |
| (66) <i>Belone belone</i> | European Garfish | 44 |
| 14. Holocentridae | | |
| (67) <i>Holocentrus ascensionis</i> | Squirrelfish | 62 |
| 15. Atherinidae | | |
| (68) <i>Melanoaenis nigrans</i> | Australian Rainbow Fish | 11 |
| 16. Serranidae | | |
| (69) <i>Centropristis furvus</i> | | 62 |
| (70) <i>Centropristis striatus</i> | Black Sea Bass | 3 |
| (71) <i>Epinephelus adscensionis</i> (<i>Epinephelus ascensionis</i>) | Rock Hind | 62 |
| (72) <i>Epinephelus guttatus</i> | Red Hind | 62 |
| (73) <i>Epinephelus morio</i> | Red Grouper | 62 |
| (74) <i>Epinephelus</i> sp. | Gray Grouper | 62 |
| (75) <i>Epinephelus</i> sp. | Queen Grouper | 62 |
| (76) <i>Epinephelus striatus</i> | Nassau Grouper | 62 |
| (77) <i>Morone americana</i> | White Perch | 62 |
| (78) <i>Morone labrax</i> | European bass | 44 |

†Dept. Path. Univ. Calif. Med. Center, Los Angeles.

TABLE IV. FISHES, AMPHIBIANS AND REPTILES IN WHICH TUBERCULOSIS HAS BEEN REPORTED (Continued)

| Species | Common Name | Reference |
|---|-----------------------------|------------|
| (79) <i>Mycteroperca bonaci</i> (<i>Epinephelus</i> var.) | Black Grouper | 62 |
| (80) <i>Mycteroperca falcata</i> (<i>Mictoperca phenax</i>) | Scamp | 62 |
| (81) <i>Roccus saxatilis</i> (<i>Roccus lineatus</i>) | Striped bass | NYA, 3, 62 |
| 17. Centrarchidae | | |
| (82) <i>Lepomis gibbosus</i> (<i>Eupomotis gibbosus</i>) | Pumpkinseed | 11 |
| (83) <i>Micropterus dolomieu</i> | Smallmouth Bass | 62 |
| 18. Percidae | | |
| (84) <i>Lucioperca lucioperca</i> (<i>Lucioperca sandra</i>) | European Pike-Perch | 23 |
| (85) <i>Perca flavescens</i> | Yellow Perch | 62 |
| 19. Carangidae | | |
| (86) <i>Trachinotus carolinus</i> | Common Pompano | 62 |
| (87) <i>Vomer setapinnis</i> | Moonfish | 62 |
| 20. Lutianidae | | |
| (88) <i>Ocyurus chrysurus</i> | Yellow-tailed snapper | 62 |
| (89) <i>Lutianus apodus</i> | Schoolmaster | 62 |
| (90) <i>Lutianus griseus</i> | Gray snapper | 3, 62 |
| (91) <i>Lutianus jocu</i> | Dog snapper | 62 |
| (92) <i>Lutianus synagris</i> | Lane (Spot) snapper | 62 |
| 21. Sciaenidae | | |
| (93) <i>Cynoscion regalis</i> | Weakfish or Gray Squeteague | 62 |
| (94) <i>Leiostomus xanthurus</i> | Spot | 62 |
| (95) <i>Micropogon undulatus</i> | Atlantic Croaker | 3, 62 |
| (96) <i>Pogonias cromis</i> | Black Sea Drum | 62 |
| 22. Pomadasyidae | | |
| (97) <i>Bathystoma</i> sp. | Red-Mouth Grunt | 62 |
| (98) <i>Plectorhynchus</i> sp. | Sweet Lip | NYA |
| (99) <i>Anisotremus surinamensis</i> | Black Margate | 62 |
| (100) <i>Anisotremus virginicus</i> | Porkfish | 62 |
| 23. Toxotidae | | |
| (101) <i>Toxotes jaculator</i> | Archerfish | NYA |
| 24. Kyphosidae | | |
| (102) <i>Kyphosus secatrix</i> | Bermuda Chub | 62 |
| 25. Sparidae | | |
| (103) <i>Archosargus probatocephalus</i> | Sheephead | 62 |
| (104) <i>Sargus sargus</i> (<i>Sargus rondeleti</i>) (<i>Diplodus sargus</i>) | Sargo | 32 |
| (105) <i>Cantharus lineatus</i> | Oldwife or Black Sea Bream | 32 |
| 26. Maenidae | | |
| (106) <i>Spicara argus</i> (<i>Smaris alcedo</i>) | Picarel Martin Pêcheur | 44 |
| 27. Scatophagidae | | |
| (107) <i>Scatophagus argus</i> | Scat | 32 |
| (108) <i>Monodactylus argenteus</i> | Silver Angelfish | 32 |
| 28. Cichlidae | | |
| (109) <i>Apistogramma ramirezi</i> | Ramirez's dwarf cichlid | 43, 45 |
| (110) <i>Cichlasoma jacetum</i> | Chanchita | 32 |
| (111) <i>Cichlasoma biocellatum</i> | Jack Dempsey | 32 |
| (112) <i>Cichlasoma festivum</i> (<i>Cichlasoma insignis</i>) | Festive Cichlid | 32 |
| (113) <i>Cichlasoma meeki</i> | Firemouth | 43, 45 |
| (114) <i>Aequidens portalegrensis</i> | Port | NYA, 2 |
| (115) <i>Aequidens curviceps</i> | Flag Cichlid | 2 |
| (116) <i>Haplochromis multicolor</i> | Egyptian Mouth-Breeder | 2, 27 |
| (117) <i>Hemichromis bimaculatus</i> | Jewelfish | 11 |
| (118) <i>Nannacara anomala</i> | Golden-Eyed Dwarf Cichlid | 2, 27 |

TABLE IV. FISHES, AMPHIBIANS AND REPTILES IN WHICH TUBERCULOSIS HAS BEEN REPORTED (Continued)

| Species | Common Name | Reference |
|--|--------------------------------------|-----------------------|
| (119) <i>Pterophyllum scalare</i> , or <i>Pterophyllum eimekei</i> | Angelfish or Scalare | NYA, 11, 32 |
| (120) <i>Symphysodon discus</i> | Disc Cichlid | NYA, 43, 45 |
| 29. Pomacentridae | | |
| (121) <i>Abudefduf saxatilis</i> (<i>Abudefduf marginalis</i>) (<i>Abudefduf mauritii</i>) | Sergeant Major | 3, 62 |
| (122) <i>Amphiprion percula</i> | Clownfish | NYA |
| (123) <i>Amphiprion laticlavius</i> | White-Saddled Clownfish | NYA |
| (124) <i>Amphiprion xanthurus</i> | Chocolate Clownfish | NYA |
| (125) <i>Amphiprion akallaopsis</i> | Skunkfish | NYA |
| (126) <i>Dascyllus auratus</i> | White-Tailed Puller | NYA, 32 |
| (127) <i>Pomacentrus leucostictus</i> (<i>Eupomacentrus leucostictus</i>) | Beau Gregory | 62 |
| (128) <i>Premnas biaculeatus</i> | Spiny Clownfish | NYA |
| 30. Chaetodontidae | | |
| (129) <i>Pomacanthus arcuatus</i> | Black Angelfish | 62 |
| (130) <i>Angelichthys isabelita</i> | Blue Angelfish (common Angelfish) | 62 |
| 31. Labridae | | |
| (131) <i>Lachnolaimus maximus</i> | Hogfish | 62 |
| (132) <i>Tautog onitis</i> | Tautog | 62 |
| 32. Acanthuridae | | |
| (133) <i>Acanthurus coeruleus</i> (<i>Teuthis coeruleus</i>) | Blue Tang | 62 |
| 33. Anabantidae | | |
| (134) <i>Anabas testudineus</i> | Climbing Perch | NYA, 27, 32 |
| (135) <i>Betta splendens</i> | Siamese Fighting Fish | NYA, 11 |
| (136) <i>Colisa lalia</i> | Dwarf Gourami | NYA, 11, 27 |
| (137) <i>Macropodus opercularis</i> | Paradisefish | NYA, 11 |
| (138) <i>Trichogaster trichopterus</i> (<i>Trichopodus trichopterus</i>) | Three-Spot (Blue) Gourami | NYA, 11, Conroy, 1963 |
| (139) <i>Trichogaster leeri</i> | Pearl Gourami | NYA, 32, Conroy, 1963 |
| (140) <i>Trichopsis vittatus</i> (<i>Ctenops vittatus</i>) | Croaking Gourami | 11 |
| 34. Triglidae | | |
| (141) <i>Prionotus carolinus</i> | Common Sea Robin | 62 |
| 35. Pleuronectidae | | |
| (142) <i>Hippoglossus hippoglossus</i> (<i>Hippoglossus vulgaris</i>) | Atlantic halibut | 29, 56 |
| 36. Bothidae | | |
| (143) <i>Lophopsetta maculata</i> | Windowpane | 62 |
| (144) <i>Paralichthys dentatus</i> | Fluke or Summer Flounder | 62 |
| 37. Monacanthidae | | |
| (145) <i>Alutera monacanthus</i> | Filefish | 62 |
| 38. Balistidae | | |
| (146) <i>Balistes vetula</i> | Queen triggerfish | 62 |
| (147) <i>Balistes carolinensis</i> | Common triggerfish | 62 |
| 39. Diodontidae | | |
| (148) <i>Chilomycterus schoepfi</i> | Spiny Boxfish or Burrfish | 62 |
| (149) <i>Diodon hystrix</i> | Porcupine fish | 62 |
| (150) <i>Sphaeroides maculatus</i> | Northern puffer | 62 |
| 40. Batrachoididae | | |
| (151) <i>Opsanus tau</i> | Northern Toadfish | 62 |
| II. Amphibia | | |
| 1. Ambystomidae | | |
| (1) <i>Siredon mexicanus</i> (= <i>Ambystoma mexicanus</i>) | Axolotl | 23, 43 |

TABLE IV. FISHES, AMPHIBIANS AND REPTILES IN WHICH TUBERCULOSIS HAS BEEN REPORTED (*Continued*)

| Species | Common Name | Reference |
|--|----------------------------|----------------|
| 2. Pseudidae | | |
| (2) <i>Pseudis paradoxa</i> | Paradox Frog | 22, 23 |
| 3. Ranidae | | |
| (3) <i>Rana catesbeiana</i> | American Bullfrog | 5 |
| (4) <i>Rana</i> spp. | European Frogs | 31, 30, 33, 50 |
| (5) <i>Rana tigrina</i> | | 26 |
| 4. Leptodactylidae | | |
| (6) <i>Leptodactylus pentadactylus</i> | Robber Frog | 13 |
| (7) <i>Pleurodema cinere</i> | | 35 |
| (8) <i>Pleurodema marmoratus</i> | | 35 |
| (9) <i>Ceratophrys americana</i> | South American Horned Frog | 26 |
| 5. Bufonidae | | |
| (10) <i>Bufo spinulosus</i> | | 35 |
| 6. Pipidae | | |
| (11) <i>Xenopus laevis</i> | South African Clawed Frog | 52 |
| III. Reptilia | | |
| 1. Chelonidae | | |
| (1) <i>Chelone corticata</i> (= <i>Caretta caretta</i>) | Loggerhead Turtle | 17 |
| 2. Trionychidae | | |
| (2) <i>Trionyx gangeticus</i> | Indian Softshell Turtle | 53 |
| (3) <i>Trionyx triungis</i> | Nile Softshell Turtle | 11, 23 |
| 3. Alligatoridae | | |
| (4) <i>Caiman sclerops</i> | Spectacled Caiman | 22, 23, 24, 53 |
| 4. Iguanidae | | |
| (5) <i>Ctenosaura multipinnis</i> (= <i>Ctenosaura acanthura</i>) | Spiny-tailed Iguana | 53 |
| 5. Lacertidae | | |
| (6) <i>Lacerta viridis</i> | Green Lizard | 9 |
| 6. Boidae | | |
| (7) <i>Boa constrictor</i> | Boa Constrictor | 18 |
| (8) <i>Python molurus</i> | Indian Rock Python | 50 |
| (9) <i>Python reticulatus</i> | Reticulated Python | 25 |
| (10) <i>Python sebae</i> | African Rock Python | 24, 53 |
| (11) <i>Python</i> sp. | | 18, 23, 26, 54 |
| (12) <i>Python spilotes</i> | | 26 |
| 7. Colubridae | | |
| (13) <i>Coluber longissimus</i> (= <i>Elaphe longissimus</i>) | Aesculapian Snake | 23, 53 |
| (14) <i>Lampropeltis getulus holbrooki</i> | Speckled King Snake | 23, 24 |
| (15) <i>Tropidonotus natrix</i> var. <i>murorum</i> (= <i>Natrix natrix</i>) | European Grass Snake | 55 |
| (16) <i>Natrix piscator</i> | Checkered Keelback | 23, 24 |
| (17) <i>Coluber catenifer</i> (= <i>Pituophis catenifer</i>) | Gopher Snake | 4 |
| (18) <i>Tarbophis fallax</i> | Cat Snake | 22 |
| (19) <i>Thamnophis sirtalis</i> | Eastern Garter Snake | 4 |
| (20) <i>Agkistrodon piscivorus</i> | | 26 |
| 8. Elapidae | | |
| (21) <i>Naja naja</i> | Cobra | NYA, 58, 59 |
| 9. Crotalidae | | |
| (22) <i>Crotalus exsul</i> (= <i>Crotalus ruber</i>) | Red Rattlesnake | 23, 53 |
| 10. Viparidae | | |
| (23) <i>Bitis arietans</i> | Puff Adder | 23, 53 |
| 11. Anguidae | | |
| (24) <i>Anguis fragilis</i> | European Slowworm | 8 |

The disease has been found in 151 species of fishes, 11 species of amphibians and 24 species of reptiles.

In the New York Aquarium, tuberculosis was present in 40 species of fishes, especially in tropical freshwater forms belonging to the families Characidae, Cyprinidae and Poeciliidae. Of special interest is the report that the acid-fast bacillus isolated from the Neon Tetra (*Hyphessobrycon innesi*) is identical with *Mycobacterium fortuitum*, a human pathogen. The disease is also found in a group of stenohaline, Pacific coral reef species commonly called clownfishes, members of the family Pomacentridae.

The pathology of tuberculosis in the fishes studied in the Aquarium is described and the taxonomy of mycobacteria from cold-blooded vertebrates is briefly discussed.

Stock cultures of mycobacteria isolated from the Neon Tetra, Blue Gourami, Archerfish, Sweet-lip (*Plectorhynchus* sp.) and from the clownfishes are maintained in the laboratory of the New York Aquarium.

REFERENCES

1. ALEXANDER, D. M.
1913. A review of piscine tuberculosis, with a description of acid-fast bacillus in the cod. Lancashire Sea Fisheries Lab., Rep. for 1912, no. 21: 43-50.
2. AMLACHER, ERWIN
1961. Taschenbuch der Fischkrankheiten. Gustav Fischer Verlag, Jena. 286 pp.
3. ARONSON, J. D.
1926. Spontaneous tuberculosis in salt water fish. J. Infect. Dis., 39: 315-320.
4. 1929. Spontaneous tuberculosis in snakes. N. sp. *Mycobacterium thamnopheos*. J. Infect. Dis., 44: 215-223.
5. 1938. Tuberculosis of cold-blooded animals. In: Tuberculosis and Leprosy, AAAS, Symposium Series, vol. 1: 80-86.
6. BAKER, J. A. & W. A. HAGAN
1942. Tuberculosis of the Mexican platyfish (*Platypoecilus maculatus*). J. Infect. Dis., 70: 248-252.
7. BATAILLON, E., DUBARD & L. TERRE
1897. Un nouveau type de tuberculose. Compt. Rend. Soc. Biol., 49: 446-449.
8. BATAILLON, E., A. MOELLER & L. TERRE
1902. Über die Identität des Bacillus des Karpfens (Bataillon, Dubard und Terre) und des Bacillus der Blindsschleuche (Moeller). Z. Tuberk., 3: 467-468.
9. BERTARELLI, E. & J. BOCCHIA
1910. Neue Untersuchungen über die Tuberkulose der Kaltblütler. Zentrbl. Bakt., Abt. I, Orig., 54: 385-393.
10. BESSE, P.
1949a. Epizootie à bacilles acid-resistants chez poissons exotique. Bull. Acad. Vet. de France, 22: 151-154.
1949b. Quelques affections à bacilles acid-resistants chez les poecilothermes vivant en aquarium. Bull. Acad. Vet. de France, 22: 425-431.
12. BOJALIL, L. F.
1959. Estudio comparativo entre *Mycobacterium marinum* y *Mycobacterium balnei*. Rev. Lat. Amer. Microbiol., 2: 169-174.
13. DARZIN, E.
1952. The epizootic of tuberculosis among the Ghias in Bahia. Acta Tuberc. Scand., 26: 170-174.
14. EARP, BRIAN J.
1959. Mycobacteria in adult salmonoid fishes returning to State hatcheries in Washington. U.S. Fish & Wildlife Serv., Sp. Sci. Rep. Fish. no. 332: 10-12.
15. EARP, B. J., C. H. ELLIS & E. J. ORDAL
1953. Kidney disease in young salmon. State of Washington, Dept. Fish., Sp. Rept., Ser. no. 1, March 1953, 74 pp.
16. FREGNAN, G. B., D. W. SMITH & H. M. RANDALL
1961. Biological and chemical studies on mycobacteria. Relationship of colony morphology to mycoside content for *Mycobacterium kansasii* and *Mycobacterium fortuitum*. J. Bact., 82: 517-527.
17. FRIEDMANN, F. F.
1903. Spontane Lungentuberkulose mit grosser Kaverne bei einer Wasserschildkröte (*Chelone corticata*). Deutsch. med. Wochenschr., 29: 25-26.
18. GIBBES, H. & E. L. SHURLY
1890. An investigation into the etiology of phthisis. Amer. J. Med. Sci., 100: 145-158.
19. GORDON, RUTH
1957. Genus 1. *Mycobacterium* Lehmann and Newmann, 1896. In: Bergey's Manual of Determinative Bacteriology, 7th Ed., pp. 695-705. Williams & Wilkins Co., Baltimore.
20. GORDON, RUTH E. & JOAN M. MIHM
1959. A comparison of four species of mycobacteria. J. Gen. Microbiol., 21: 736-748.

21. GORDON, RUTH E. & MILDRED M. SMITH
1955. Rapidly growing acid-fast bacteria. II. Species description of *Mycobacterium fortuitum* Cruz. J. Bact., 69: 502-507.
22. Griffith, A. S.
1928. Tuberculosis in captive wild animals. J. Hyg., 28: 198-218.
23. 1930. Tuberculosis in cold-blooded animals. In: A System of Bacteriology in Relation to Medicine. Med. Res. Council. His Majesty's Stationary Office, London, 5: 326-332.
24. 1941. The susceptibility of the water (or grass) snake (*Tropidonotus natrix*) to the avian tubercle bacillus and to reptilian strains of acid-fast bacilli. J. Hyg., 41: 284-288.
25. HANSEMANN, VON
1903. Ueber saurefest Bacillen bei *Python reticularis*. Zentrbl. Bakt., Abt. I, Orig., 34: 212-213.
26. IPPEN, R.
1962. Die Spontane Tuberkulose bei Kaltblütern. Nord. Vet.-Med., 14, suppl. 1: 183-192.
27. JAHNEL, J.
1940. Die Fischtuberkulose. Wochenschr. f. Aquarien und Terrarienkunde, 37: 317-321.
28. JOHNSTONE, J.
1913. Diseased conditions of fishes. Lancashire Sea Fisheries Lab., Rept. no. 21, 1912: 2-42; see also Proc. & Trans. Liverpool Biol. Soc., 27: 196-201, 1912-1913.
29. 1927. Diseased conditions of fishes. Lancashire Sea Fisheries Lab., Rept. no. 35, 1926: 162-167.
30. KÜSTER, E.
1905. Ueber Kaltblütertuberkulose. Much. Med. Wochenschr., 52: 57-58.
31. 1928. Die Kaltblütertuberkulose. In: Kolle und Wassermann's Handbuch der Pathogenen Mikroorganismen, 5: 746-774.
32. LEDERER, GUSTAV
1943. Fishtuberkulose. Der Zoologische Garten (N.F.), 15: 107-121.
33. LICHTENSTEIN, S.
1943. Ein Fall von Spontanen Froschtuberkulose. Zentrbl. Bakt. Abt. I, Orig., 85: 249.
34. LINELL, F. & A. NORDEN
1954. *Mycobacterium balnei*. A new acid-fast bacillus occurring in swimming pools and capable of producing skin lesions in humans. Acta. Tuberc. Scand. (suppl.), 33: 1-85.
35. MACHICAO, N. & E. LA PLACA
1954. Lepralike granulomas in frogs. Lab. Invest., 3: 219-227.
36. McMILLEN, SHIRLEY
1960. *Mycobacterium fortuitum*: a mutant of *Mycobacterium marinum*. Bact. Proc., p. 79.
37. McMILLEN, SHIRLEY & DANIEL S. KISHNER
1959. *Mycobacterium marinum* Aronson (1926). Bact. Proc., p. 31.
38. NIGRELLI, ROSS F.
1953. Causes of diseases and death of fishes in captivity. Zoologica, 28: 203-216.
39. 1953. Two diseases of the Neon Tetra, *Hyphessobrycon innesi*. Aquarium Journal (San Francisco), 24: 203-208.
40. PARISOT, THOMAS J.
1958. Tuberculosis in fish. Bact. Rev., 22: 240-245.
41. PARISOT, THOMAS J. & EDWARD M. WOOD
1960. A comparative study of the causative agent of the mycobacterial disease of salmonoid fishes. II. A description of the histopathology of the disease in Chinook salmon (*Oncorhynchus tshawytscha*) and a comparison of the staining characteristics of the fish disease with leprosy and human tuberculosis. Amer. Rev. Respiratory Dis., 82: 212-222.
42. REED, G. B.
1948. Mycobacteriaceae. In: Bergey's Manual of Determinative Bacteriology, 6th Ed., pp. 875-891. Williams & Wilkins Co., Baltimore.
43. REICHENBACH-KLINKE, H. H.
1955a. Die Fischtuberkulose. Die Aq.-und-Terr.-Zeitschr., No. 1: 12-17.
44. 1955b. Beobachtungen über Meeresfisch-Tuberkulose. Pubbl. Stazione Zoologica, Napoli, 26: 55-62.
45. 1957. Krankheiten der Aquarienfische. Alfred Kernen Verlag, Stuttgart. 215 pp.
46. ROSS, A. JOHN
1959. Mycobacteria in adult salmonoid fishes returning to federal hatcheries in Washington, Oregon and California. U.S. Fish & Wildlife Serv., Sp. Sci. Rept. Fish., no. 332: 1-9.
47. 1960. *Mycobacterium salmoniphilum*, sp. nov., from salmonoid fishes. Amer. Rev. Respiratory Dis., 81: 241-250.
48. ROSS, A. JOHN & F. P. BRANCATO
1959. *Mycobacterium fortuitum* Cruz from the tropical fish *Hyphessobrycon innesi*. J. Bact., 78: 392-395.

49. ROSS, A. J. & H. E. JOHNSON
1962. Studies of transmission of mycobacterial infections in Chinook salmon. *The Progressive Fish-Culturist*, 24: 147-149.
50. RUPPRECHT, J.
1904. Über Säurefestes nebst Beschreibung eines Falles von spontaner Froschtuberkulose. Inaug. Diss. Freiburg; also in: *Jber. Lehre pathog. Mikroorganism (1904)*, 20: 591.
51. SCHREITMÜLLER, W. & G. LEDERER
1929. *Fischkrankheiten. Das Aquarium*, 3: 26.
52. SCHWABACHER, H.
1959. A strain of *Mycobacterium* isolated from skin lesions of a cold-blooded animal, *Xenopus laevis*, and its relation to a typical bacilli occurring in man. *J. Hyg.*, 57: 57-67.
53. SCOTT, H. H.
1930. Tuberculosis in man and lower animals. *Med. Res. Council (Great Britain)*, Sp. Rept., Ser., 149: 1-270.
54. SHATTOCK, S. G.
1902. Tuberculosis of the esophagus in a *Python*. *Tran. Path. Soc., London*, 53: 430-440.
55. SIBLEY, W. K.
1899. Ueber Tuberkulose bei Wirbelthieren. *Virchow's Arch.*, 116: 105-115.
56. SUTHERLAND, P. L.
1922. A tuberculosis-like disease in a salt-water fish (halibut) associated with the presence of an acid-fast tubercle-like bacillus. *J. Path. & Bact.*, 25: 31-35.
57. SWIFT, SHELDON & HAROLD COHEN
1962. Granuloma of the skin due to *Mycobacterium balnei* after abrasions from a fish tank. *New England J. Med.*, 267: 1244-1246.
58. VOGEL, HENRY
1956. Metabolic and serologic study of acid-fast bacteria from fishes and other cold-blooded vertebrates. Doctoral Dissertation, Dept. Biol., N. Y. Univ. Graduate Sch. Arts and Sci.
59. 1958. Mycobacteria from cold-blooded animals. *Amer. Rev. Tuberculosis*, 77: 823-838.
60. 1959. A metabolic study of acid-fast bacteria from cold-blooded animals. *J. Infect. Dis.*, 104: 28-37.
61. WESTGARD, RICHARD L.
1959. A procedure for the detection of acid-fast bacteria in fish: A comparative study of digest-concentrate vs. smear technique. *U. S. Fish & Wildlife Serv., Sp. Rept. Fisheries*, no. 332: 13-17.
62. WINSOR, HENRY
1946. Cold-blooded tuberculosis from the Fairmount Park Aquarium, Philadelphia. *Proc. Penna. Acad. Sci.*, 20: 43-46.
63. WOOD, JAMES W.
1959. A survey of tuberculosis in Pacific salmon and steelhead trout in Oregon streams in 1957. *U. S. Fish & Wildlife Serv., Sp. Sci. Rept. Fish.*, no. 332: 18-34.
64. WOOD, JAMES W. & ERLING J. ORDAL
1958. Tuberculosis in Pacific salmon and steelhead trout. *Fish. Comm. of Oregon, Portland, Contr.* no. 25: 1-38.

EXPLANATION OF THE PLATES

PLATE I

- FIG. 1. Neon Tetra infected with tuberculosis. Note various degrees of emaciation. A yellowish discoloration of the usually brilliant red markings on the lateral-posterior area is an external manifestation of the disease. Slightly larger than natural size. Photo by S. C. Dunton, New York Zoological Society.
- FIG. 2. Typical acid-fast organisms in smear from discolored area of the skin of the Neon Tetra. Ziehl-Neelsen's stain. 650 \times .

PLATE II

- FIG. 3. Section through the body wall and muscle of the Neon Tetra showing numerous solitary and coalescent tubercles. Hematoxylin-eosin. 100 \times .
- FIG. 4. Section through the body of an infected Neon Tetra showing a characteristic condition of tuberculosis in tropical freshwater fish. Upper left is the liver; the intestine is shown slightly to right of center; in between is the mesentery in which the diffuse exocrine pancreas is present. Hematoxylin-eosin. 50 \times .

PLATE III

- FIG. 5. Stained smear of the internal organs of infected Neon Tetra showing the typical acid-fast mycobacteria, some of which occur within macrophages. Ziehl-Neelsen's stain. 650 \times .

- FIG. 6. Electrophotomicrograph of acid-fast bacillus from a culture of the Neon Tetra Strain-N. 16,300 \times .

PLATE IV

- FIG. 7. Liver of the cyprinodon *Cynolebias* sp. showing the epitheloid-like tubercles in the parenchyma. Hematoxylin-eosin. 450 \times .
- FIG. 8. Higher magnification of the central area of a tubercle shown in fig. 7. Note the typical acid-fast bacilli. Ziehl-Neelsen. 650 \times .

PLATE V

- FIG. 9. Three-spot or blue gourami with lepromatous-like macular and necrotic skin and fin lesions. $\frac{1}{2}$ natural size.
- FIG. 10. Hard tubercles ("pearl bodies") in liver and other organs of the Climbing Perch, *Anabas testudineus*. Slightly larger than natural size.

PLATE VI

- FIG. 11. Section in region of the kidney of a goldfish showing coalescent and some degree of caseation of the tubercles. Note lymphocytic infiltration and fibrous development. Hematoxylin-eosin. 300 \times .
- FIG. 12. Luxuriant subculture of *Mycobacterium*, Neon Tetra Strain-H, originally isolated from the kidney; grown on Dorset's egg agar at 37°C. A subculture of this strain has been identified as *Mycobacterium fortuitum* by Ross & Brancato (1959).